



Case report

Gunshot wound in skeletonised human remains with partial adipocere formation

Faridah Mohd Nor MD, PhD Associate Professor^{a,*},
Srijit Das MBBS, MS Associate Professor^b

^a Department of Pathology, Universiti Kebangsaan Malaysia Medical Centre, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, 56000 Kuala Lumpur, Malaysia

^b Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

ARTICLE INFO

Article history:

Received 21 October 2010

Accepted 20 July 2011

Available online 9 August 2011

Keywords:

Exhumation

Skeletonised

Human

Remains

Bullet

Adipocere

Forensic

Autopsy

ABSTRACT

We present a case of skeletonised human remains. In the present case report, a body was exhumed from the ground above a cemetery. On exhumation, the body was partially-skeletonised with adipocere formation on the upper part of the body. Autopsy of the body showed two bullets in the right thigh muscle and lumbar vertebrae between L4 and L5. Postmortem changes and destruction of soft tissue made it impossible to determine direction of fire through the body, even in a careful complete autopsy.

© 2011 Elsevier Ltd and Faculty of Forensic and Legal Medicine. All rights reserved.

1. Introduction

Exhumation can be defined as the act of digging something from the burial ground.¹ It is performed only for special reasons such as moving the body to a different location or to be buried closely with family members. Taphonomic effects of erosion on deliberately buried bodies have been studied. In the case of wind erosion, the bone becomes very poorly preserved while, water erosion may cause both the remains and the deposits to be lost at the same time, although the bones may be well preserved. This observation is particularly useful for the investigation of clandestine burial and time since deposition.² Ground penetrating radar (GPR) was used to detect cadavers buried in different types of soil, and it was found that it may be difficult to detect cadavers soon after they are skeletonised as the surrounding area may not provide a strong contrasting area to be detected by GPR. Also, bodies buried at deeper depths may not give good detection as they showed reduced decomposition rates.³ Alternatively, odour analysis of decomposing buried human remains may be used to detect human remains in burial grounds.⁴

Adipocere is often being found in buried bodies. Adipocere is yellowish, cheesy product which has an ammoniacal odour. There are factors in the surrounding soil either accelerate or slow the

formation of adipocere in buried bodies.⁵ It was documented that soil conditions particularly its water content is a crucial factor in enhancing the formation of adipocere,⁶ while decomposition of soft tissue will stimulate changes in the soil microenvironment.⁷ In addition, the presence of coffins slows down the formation of adipocere, but clothings enhance its formation.⁸ Literature has depicted that the time required for adipocere formation varies from three days to five years.⁹ Nevertheless, aeration of adipocere in corpses disappear within several years.¹⁰

Identification in fragmented human bones can be difficult due to commingled human and animal remains. The process of documentation and handling of multiple fragmented human bones have been positively identified in death cases in Germany especially during world war II.¹¹ Besides having morphologic means of identification, mtDNA of skeletal remains may be useful even in human remains, buried centuries ago.¹² It has been documented that postmortem radiographic examination is useful in determining gunshot injuries and bullet locations both in human and in animals.^{13,14} Here, we describe a case of a buried body with gunshot wounds, in which computerized tomography (CT) and magnetic resonance imaging (MRI) were used to localize the bullets.

2. Case report

In this case report, we describe a body, which was buried in a clandestine burial above an old grave. Identification of the grave

* Corresponding author. Tel.: +60 006 03 91455368; fax: +60 006 03 91737340.
E-mail addresses: mnfaridah@yahoo.com, mnfaridah@gmail.com (F. Mohd Nor).

was done by the cemetery authorities and family members of the deceased. This was necessary to respect the body and to prevent further disturbance. The grave yard was closed by the police from possible interference by the public or media. The criminal suspects were brought in to identify the grave before the magistrate and the pathologist. Protective and disposable masks, gowns, and gloves were worn by the personnel to protect themselves from possible transmission of disease.

Exhumation of the grave showed that the deceased's head was buried under a cement block, which was 43 cm below the ground surface. The body was clad in short-sleeved shirt and trousers, and was in left lateral position with all four limbs flexed. Some adipocere formation was found on upper part of the body with partially-skeletonised lower body. The skeletal remains were dug in one piece along with some soil into a body bag. This was done to avoid loss of evidence and to prevent unnecessary movement of the skeleton. Collection of soil samples from the grave was not performed, as there was no suspicion of drug or heavy metal poisoning. In sudden death, when other specimens are unavailable due to degradation, bone marrow can be useful specimen for detection of drug in drug poisoning.¹⁵ The body was buried in a grave measuring 2.3 metres \times 85 cm 85 cm in length, width and depth, respectively.

In one study, TraumaSCAN-Web, a Java based system was used to assess thoraco-abdominal penetrating trauma from gunshot and stab wounds, and this system is based on geometric calculations on anatomic involvement in injury.¹⁶ However, in this case, computerized tomography (CT) and magnetic resonance imaging (MRI) were used to assess the skeletal remains prior to autopsy. It showed presence of two bullets between L4 and L5 (Fig. 1) vertebrae and distal right thigh. The bullets (copper-jacketed) retrieved from the vertebrae and right thigh measured 1.3 by 0.4 cm and 1.2 by 0.4 cm, respectively. The bullets were sent to a ballistics laboratory as evidence, where they were confirmed to be 9-mm copper-jacketed bullets.

Postmortem examination in exhumation cases is basically the same as in fresh body, but may require some modifications where necessary.¹⁷ A lot of information may be gained from exhumed cases, for instance soft tissues may survive between 8 years to about 16 years of burial time. This is particularly applicable for burials in cold temperature, in which changes of injuries may still be visible.^{18,19} Nevertheless, in this case the soft tissue was completely disintegrated. There was no skin or soft tissues to document the bullet pathway in the body. The soft tissues had

disintegrated due to postmortem changes. The body was partially-skeletonised on lower part of the body, in which several tattoos were present on the abdomen, back and anterior left thigh. This was particularly useful for identification purposes. In addition, two bullets were retrieved from the right thigh and between the lumbar vertebrae, L4 and L5. The vertebrae were not fractured, but the medial condyle of left femur showed a bullet path with surrounding contusions.

Anthropologic assessment showed an incomplete skeletal inventory with several extremities bones missing. The coccyx bone showed a right lateral deviation, which provided a unique identification feature (Fig. 2a). Additionally, intra-medullary nailing was present in the right femur with the end of the nail abutting the bone proximally (Fig. 2b). There was evidence of partially fused spiral fracture with spurs and callus formation of the middle third of right femur. There were some peculiar greenish stains in the distal right thigh, which was consistent with that of a copper-jacketed bullet. A bullet was retrieved from the medial condyle of right femur with surrounding contusion (Fig. 2c and d).

Further, the stature was determined by using mean values of methods by 'Dupertuis and Hadden's' and 'Trotter and Gleser'. Racial affiliation was determined by using the average of several indices such as cephalic, brachial, crural, inter-membral and intra-membral indices. The age was estimated by taking mean values from examination of cranial suture closure, pubic symphyseal surfaces and rib bone. Sex was assessed by examining pelvis, femur, humerus, skull, long bone, sternum, mandible and scapula. Based on anthropologic assessment, the deceased was European male with mean stature of 178.07 cm and estimated age of 41 years old.

3. Discussion

According to the suspect, the deceased had received two shots in the right knee and abdomen. It may be argued that for the bullet to reach lumbar vertebrae, two possibilities are anticipated. The shot could have been discharged to the abdomen or it could be a shot directly to the back. If a shot to the abdomen is anticipated, the bullet would have lost its kinetic energy upon traversing the peritoneum and internal organs in order to reach the vertebrae. However, the presence of advanced decomposition with absence of soft tissue has made it very difficult to determine direction of fire in this case. On the contrary, it could also be a shot direct to the back, which may or may not show fracture of the vertebrae. Studies have shown that penetrating bullet to the back did not cause fracture of the surrounding vertebrae, but dissipates thermal and kinetic energy into the surrounding tissues.²⁰ In this case, the lumbar vertebrae were not fractured. Hence, the allegations made by the suspects could either be true or false.

As per the police, the deceased was a criminal suspect, who had involved in car thefts around the country. He was kidnapped by a group of gangsters, who had shot him in the right knee. A CCTV at a hotel had successfully captured his picture, who was seen limping on his right leg. He received a fatal shot to the abdomen and died. He was buried in a clandestine grave with the help of his accomplices. The body was later dug by animals and re-buried, during which cement mixture was poured to the head in order to cover it completely.

In this case, the deceased was shot and buried in a clandestine burial. Because the deceased was a criminal, the license for exhumation was issued by the magistrate court to gain new forensic evidence. The identification of the deceased was made by the suspects based on the identification of the burial place, the deceased and his personal effects. In brief, the circumstantial evidence well corroborated to the accounts given by the suspects.




Fig. 1. Photograph of CT scan and MRI demonstrating a bullet between two lumbar vertebrae, L4 and L5.



Fig. 2. a. Photograph showing right lateral deviation of coccyx (identification). b. Photograph showing intra-medullary nailing of proximal right femur. c. Photograph showing contused bullet path on antero-medial aspect of distal right femur. d. Photograph showing bullet path (close-up).

The presence of tattoos on the body and intra-medullary nailing (IMN) in the right femur had further confirmed identification of the deceased. The location of bullets had also corroborated to the allegations made by the suspects. As per identification, the deceased was an Indian male, which was consistent with Caucasoid features based on anthropologic assessment.

Advanced decomposition with partial skeletalisation and adipocere formation of the body indicated that the individual died between a time period ranging from few weeks to months. According to the suspects, the body had been buried for about four months, which was largely consistent with time of death and his reported missing. Adipocere formation on upper part of the body was contributed by the presence of damp soil around the body. Even in a careful complete autopsy, cases have shown that it can be very difficult to show direction of fire in decomposed bodies.²¹ The absence of skin and soft tissues further reduces the chances of finding direction of fire through the body. One case study reported that a transperitoneal shot to the abdomen had caused a bullet to be lodged between L5 and S1 vertebrae.²² In this case, a transperitoneal shot to the abdomen had caused a bullet to be lodged between L4 and L5 vertebrae. Studies also showed that injuries to the lower lumbar spine in gunshot wounds were infrequent and involved the transfer of high amounts of energy. Of the injuries involving lumbar spine, pedicle fractures were among the least common, and those involving bilateral pedicles were even rare.²³ This is because the lower lumbar spine being well protected from traumatic injury by its strong muscular and ligamentous attachments to the pelvis and axial skeleton. Injuries to the area involve the transfer of high amounts of energy.²³

The distinction between homicide, suicide and accident can sometimes be extremely difficult and a final conclusion can only be reached after a full police investigation. Nevertheless, multiple firearm wounds strongly suggest homicide.²⁴ Although, circumstantial evidence and postmortem findings showed corroborative evidence, the cause and manner of death in this case was difficult

to determine, and may remain unascertained until proven otherwise.

Acknowledgements

The author thank the doctors in Universiti Kebangsaan Malaysia Medical Centre and the police in Cheras area of Kuala Lumpur for their assistance in the investigation of the case.

Conflict of interest

None.

Funding

No funding was received.

Ethical approval

It was not necessary.

References

1. The free dictionary by Farlex, <http://www.thefreedictionary.com/exhumation>; 2010. Last accessed on 21.10.2010.
2. Littleton J. Taphonomic effects of erosion on deliberately buried bodies. *J Arch Sci* 2000;**27**:5–18.
3. Schultz JJ. Sequential monitoring of burials containing small pig cadavers using ground penetrating radar. *J Forensic Sci* 2008;**53**(2):279–87.
4. Vass AA, Smith RR, Thompson CV, Burnett MN, Dulgerian N, Eckenrode BA. Odor analysis of decomposing buried human remains. *J Forensic Sci* 2008;**53**(2):384–91.
5. Forbes SL, Stuart BH, Dent BB. The effect of the method of burial on adipocere formation. *Forensic Sci Int* 2005;**154**:44–52.
6. Durães N, Cortez D, Algarra M, Sánchez FG, Rodríguez- Borges JE, Bobos I, et al. Comparison of adipocere formation in four soil types of the Porto (Portugal) district. *Forensic Sci Int* 2010;**195**:168e.1–168e.6.
7. Wilson AS, Janaway RC, Holland AD, Dodson HI, Baran E, Pollard AM, et al. Modelling the buried human body environment in upland climes using three contrasting filed sites. *Forensic Sci Int* 2007;**169**:6–18.
8. Forbes SL, Dent BB, Stuart BH. The effect of soil type on adipocere formation. *Forensic Sci Int* 2005;**154**:35–43.
9. Mohan Kumar TS, Monteiro FNP, Bhagavath P, Bakkannavar SM. Early adipocere formation: a case report and review of literature. *J Forensic Leg Med* 2009;**16**:475–7.

10. Fründ H-C, Schoenen D. Quantification of adipocere degradation with and without access to oxygen and to the living soil. *Forensic Sci Int* 2009;**188**:18–22.
11. Hollmann T, Byard RW, Tsokos M. The processing of skeletonized human remains found in Berlin, Germany (Original communication). *J Forensic Leg Med* 2008;**15**:420–5.
12. Salerno A, Lago G, Berti A, Calì F, Mirisola M, Carta P, et al. mt DNA analysis of the human remains buried in the sarcophagus of Federico II. *J Cult Herit* 2005;**6**:313–9.
13. Thali MJ, Yen K, Vock P, Ozdoba C, Kneubuehl BP, Sonnenschein M, et al. Image-guided virtual autopsy findings of gunshot victims performed with multi-slice computed tomography (MSCT) and magnetic resonance imaging (MRI) and subsequent correlation between radiology and autopsy findings. *Forensic Sci Int* 2003;**138**:8–16.
14. Heng HG, Teoh WT, Sheikh-Omar AR. Postmortem abdominal radiographic findings in feline cadavers. *Vet Radio Ultrasound* 2008;**49**(1):26–9.
15. Cengiz S, Ulukan Ö, Ates I, Tugcu H. Determination of morphine in postmortem rabbit bone marrow and comparison with blood morphine concentrations. *Forensic Sci Int* 2006;**156**:91–4.
16. Ogunyemi O. Methods of reasoning from geometry about anatomic structures injured by penetrating trauma. *J Biomed Inform* 2006;**39**:389–400.
17. Enotes — Exhumation, <http://www.enotes.com/forensic-science/exhumation>; 2010. Last accessed on 21.10.2010.
18. MacAulay LE, Barr DG, Strongman DB. Effects of decomposition on gunshot wound characteristics: under cold temperatures with no insect activity. *J Forensic Sci* 2009;**54**(2):448–51.
19. Breitmeier D, Graefe-Kirci U, Albrecht K, Weber M, Tröger HD, Kleemann WJ. Evaluation of the correlation between time corpses spent in in-ground graves and findings at exhumation. *Forensic Sci Int* 2005;**154**:218–23.
20. Ben-Galim P, Reitman CA. Intrathecal migratory foreign body without neurological deficit after a gunshot wound. *Spine J* 2008;**8**:404–7.
21. Kobayashi M, Sakurada K, Nakajima M, Iwase H, Hatanaka K, Matsuda Y, et al. A 'keyhole' gunshot wound in an adipocere case (Brief communication). *Leg Med* 1999;**1**:170–3.
22. Tumialán LM, Walkup RR, Gupta SK. Minimally invasive retrieval of a bullet from the L5-S1 neural foramina after transperitoneal gunshot wound (Technical reviews). *Spine J* 2009;**9**:169–73.
23. Chin KR, Boselli K, Cairone S. Lag screw fixation of remote bilateral pedicle fractures of the fourth and fifth lumbar vertebrae after a single gunshot wound: a case report and technical pearl. *Spine J* 2010;**10**:136–40.
24. Desinan L, Mazzalo GM. Gunshot fatalities: suicide, homicide or accident? A series of 48 cases. *Forensic Sci Int* 2005;**147S**:S37–40.